arXiv:1212.2706v1 [cond-mat.stat-mech] 12 Dec 2012

Stochastic Echo Phenomena in Nonequilibrium Systems

Jae Dong Noh[1, 2]
[1]Department of Physics, University of Seoul, Seoul 130-743, Korea
[2]School of Physics, Korea Institute for Advanced Study, Seoul 130-722, Korea

(Dated: December 13, 2012)

A thermodynamic system is driven into a nonequilibrium condition when a time-dependent force or a nonconservative force represented by a protocol $\lambda(t)$ is applied. Such a system is time irreversible in the sense that the motion under a protocol $\lambda(0 \le t \le \tau)$ and the time-reversed motion under $\lambda(\tau - t)$ are not equally probable. We investigate a stochastic echo phenomenon in which a system is controlled by an optimal protocol to display the time-reversed motion. The optimal protocol is derived for a Langevin system. For the time-dependent driving case, simple harmonic oscillator systems are found to display the perfect stochastic echo phenomenon. We also show that any system driven by a time-independent nonconservative force has a dual system displaying the perfect stochastic echo phenomenon. Our study provides a novel perspective on the time-irreversibility of nonequilibrium systems.

PACS numbers: 05.70.Ln, 02.50.-r, 05.40.-a

Introduction – The Loschmidt echo is the characteristic feature of time reversible systems. A classical system returns to its original state when the velocity of all particles is reversed. A quantum mechanical system evolving under the Hamiltonian H and subsequently under $-H$ returns to the initial state. An echo phenomenon also occurs in a driven macroscopic system. One of the most popular examples is a drop of dye placed in a highly viscous fluid in the gap between two concentric cylinder [1]. When the inner cylinder is rotated, the drop spreads apparently uniformly in the fluid. Surprisingly, the drop reappears when the rotation is reversed. A recent experiment reports a transition between the reversible and irreversible regimes as a strength of the driving is varied [2], although it is controversial whether it is a sharp phase transition [3] or a smooth crossover [4].

An equilibrium system with Langevin dynamics obeys the detailed balance condition [5]

$$p_{eq}(\boldsymbol{x}_i) P(\boldsymbol{x}_f, t_f | \boldsymbol{x}_i, t_i) = p_{eq}(\boldsymbol{x}_f) P(\boldsymbol{x}_i, t_i | \boldsymbol{x}_f, t_f) \ , \quad (1)$$

where p_{eq} denotes the equilibrium distribution function and $P(\boldsymbol{x}', t' | \boldsymbol{x}, t)$ denotes the transition probability from $\boldsymbol{x}$ at time t to $\boldsymbol{x}'$ at time t'. The detailed balance implies that the motion along a path and the time-reversed motion are equally probable. Hence, an equilibrium system displays an echo phenomenon in the probabilistic sense.

When a thermodynamic system is exerted by a time-dependent force or a nonconservative force, it is driven into a nonequilibrium condition [6–8]. The nonequilibrium driving is represented by a protocol $\lambda(t)$, which may depend on time or not. A nonequilibrium system does not obey the detailed balance. So a motion along a path under a protocol $\lambda(0 \le t \le \tau)$ and its time-revered motion under a time-reversed protocol $\lambda^R(t) \equiv \lambda(\tau - t)$ are not equally probable. In fact, the difference in the statistical weights between them is directly related to the entropy increase, which is the basis for the various nonequilibrium fluctuation theorems [6–14].

The time irreversibility raises a question about the extent to which stochastic motions of a nonequilibrium system can be reversed in time. The absence of the detailed balance and the entropy increase indicates that a time-reversed protocol does not lead to an echo phenomenon. However, it is little known under which protocol the time-reversed motions are simulated best. We will call the controlled time-reversed motion a stochastic echo phenomenon (SEP). In this paper, we derive the condition for the optimal protocol for the SEP in nonequilibrium systems with Langevin dynamics. We begin with a review on stochastic thermodynamics for nonequilibrium Langevin systems. Then, the optimal condition for the SEP will be derived. The SEP will be further investigated in systems with time-dependent forces and with time-independent nonconservative forces. We conclude the paper with summary and discussions.

Langevin dynamics – We concentrate on the overdamped dynamics of a system of coordinate $\boldsymbol{x} = (x_1, \cdots, x_N)$ exerted by a force $\boldsymbol{f}_{\lambda(t)}(\boldsymbol{x})$. The force depends on a protocol $\lambda(t)$ representing a time-dependent force or a time-independent nonconservative force. The Langevin equation reads [5]

$$\dot{\boldsymbol{x}}(t) = \boldsymbol{f}_{\lambda(t)}(\boldsymbol{x}(t)) + \sqrt{2T} \ \boldsymbol{\eta}(t) \ , \quad (2)$$

where T is the temperature of the heat reservoir and $\boldsymbol{\eta}(t)$ is the δ-correlated thermal noise with $\langle \eta_i(t) \rangle = 0$ and $\langle \eta_i(t) \eta_j(t') \rangle = \delta_{ij} \delta(t - t')$. The damping coefficient and the Boltzmann constant are set to 1.

The initial probability density for $\boldsymbol{x}(0)$ will be denoted by $p_i(\boldsymbol{x}(0))$. Then the dynamics determines the probability density $p_f(\boldsymbol{x}(\tau))$ for $\boldsymbol{x}(\tau)$ at time $t = \tau$ and the probability density functional for a path $\boldsymbol{x}(t)$ in the interval $0 \le t \le \tau$. It is given by [15, 16]

$$\mathcal{P}_\lambda[\boldsymbol{x}] = p_i(\boldsymbol{x}(0)) e^{-\mathcal{L}[\boldsymbol{x}; \boldsymbol{f}_\lambda]} \quad (3)$$

with the action

$$\mathcal{L}[\boldsymbol{x};\boldsymbol{f}_\lambda] = \int_0^\tau dt \left\{ \frac{[\dot{\boldsymbol{x}} - \boldsymbol{f}_\lambda(\boldsymbol{x})]^2}{4T} + \frac{1}{2}\boldsymbol{\nabla}_{\boldsymbol{x}} \cdot \boldsymbol{f}_\lambda(\boldsymbol{x}) \right\} . \tag{4}$$

For convenience, we adopt the Stratonovich convention for the stochastic integral throughout the paper [5].

Suppose that the system follows a path $\boldsymbol{x}(t)$ under a protocol $\lambda(t)$. The time-reversibility and the total entropy change are related through the relation [6, 8, 10]

$$\Delta\mathcal{S}_{tot}[\boldsymbol{x}] = \ln\left[\frac{\mathcal{P}_\lambda[\boldsymbol{x}]}{\mathcal{P}^R_{\lambda^R}[\boldsymbol{x}]}\right] , \tag{5}$$

where $\mathcal{P}^R_{\lambda^R}[\boldsymbol{x}] \equiv \mathcal{P}_{\lambda^R}[\boldsymbol{x}^R]$ denotes the probability density functional for a time-reversed path $\boldsymbol{x}^R(t) \equiv \boldsymbol{x}(\tau - t)$ under a protocol $\lambda^R(t) \equiv \lambda(\tau - t)$. It can be decomposed as $\Delta\mathcal{S}_{tot}[\boldsymbol{x}] = \Delta\mathcal{S}_{sys}[\boldsymbol{x}] + \Delta\mathcal{S}_{res}[\boldsymbol{x}]$, where

$$\Delta\mathcal{S}_{sys}[\boldsymbol{x}] = -\ln\left[\frac{p_f(\boldsymbol{x}(\tau))}{p_i(\boldsymbol{x}(0))}\right] \tag{6}$$

is the change in the system entropy and $\Delta\mathcal{S}_{res}[\boldsymbol{x}] = \mathcal{Q}[\boldsymbol{x}]/T$ is the change in the heat bath entropy with the heat

$$\mathcal{Q}[\boldsymbol{x}] = \int_0^\tau dt\ \dot{\boldsymbol{x}}(t)\cdot \boldsymbol{f}_{\lambda(t)}(\boldsymbol{x}(t)) . \tag{7}$$

The quantity $-\mathcal{Q}[\boldsymbol{x}]$ is equal to the work done by the damping force and the random force. So, $\mathcal{Q}[\boldsymbol{x}]$ is the same as the heat dissipated into the heat bath.

It is noteworthy that the average value of $\Delta\mathcal{S}_{tot}$ is the relative entropy (or Kullback-Leibler divergence) of $\mathcal{P}_\lambda[\boldsymbol{x}]$ with respect to $\mathcal{P}^R_{\lambda^R}[\boldsymbol{x}]$ [17]:

$$\langle \Delta\mathcal{S}_{tot}\rangle_{\mathcal{P}_\lambda} = D\left(\mathcal{P}_\lambda \| \mathcal{P}^R_{\lambda^R}\right) . \tag{8}$$

The relative entropy of $p(x)$ with respect to $q(x)$ is defined as $D(p\|q) = \langle \ln \frac{p(x)}{q(x)}\rangle_p = \int dx\ p(x)\ln\frac{p(x)}{q(x)}$. It is nonnegative for any probability distributions p and q with $D(p\|q) = 0$ if and only if $p(x) = q(x)$ almost everywhere. So the entropy never decreases and the echo phenomenon fails with a time-reversed protocol λ^R.

Stochastic echo phenomenon – In order to find the optimal protocol for the SEP, we introduce an imaginary system subject to a force $\boldsymbol{f}_{\kappa(t)}(\boldsymbol{x})$ with a protocol $\kappa(t)$. The initial state of this system is taken to be the final state of the original system with the force $\boldsymbol{f}_\lambda$. So the probability density functional for a path $\boldsymbol{x}(t)$ in the interval $0 \le t \le \tau$ is given by $\mathcal{P}_{\kappa(t)}[\boldsymbol{x}(t)] = p_f(\boldsymbol{x}(0))e^{-\mathcal{L}[\boldsymbol{x};\boldsymbol{f}_\kappa]}$.

The optimal protocol can be obtained by minimizing the relative entropy

$$\mathbb{S}[\kappa;\lambda] = D\left(\mathcal{P}_\lambda \| \mathcal{P}^R_\kappa\right) \tag{9}$$

with respect to κ. The relative entropy is useful since it is nonnegative and zero only for the *perfect* SEP with $\mathcal{P}_\lambda[\boldsymbol{x}] = \mathcal{P}^R_\kappa[\boldsymbol{x}] = \mathcal{P}_\kappa[\boldsymbol{x}^R]$. It reduces to the average value of the total entropy change for the special choice $\kappa = \lambda^R$. Consequently, it also provides a bound for the total entropy change, $\langle \Delta S_{tot}\rangle = \mathbb{S}[\lambda^R;\lambda] \ge \min_\kappa \mathbb{S}[\kappa;\lambda] \ge 0$.

Before proceeding, we remark that the optimal protocol has also been considered in a different context. Suppose that a variation of an external parameter from λ_i at time $t = 0$ to λ_f at time $t = \tau$. One may ask a question about the optimal control of $\lambda(t)$ which yields the minimum entropy production [18, 19]. Note that this optimal protocol for the minimum entropy production is different from the optimal protocol for the SEP.

For a notational simplicity, we introduce $\mu(t) \equiv \kappa^R(t) = \kappa(\tau - t)$. The relative entropy is rewritten as

$$\mathbb{S}[\kappa;\lambda] = \left\langle -\ln\left[\frac{p_f(\boldsymbol{x}(\tau))}{p_i(\boldsymbol{x}(0))}\right]\right\rangle_{\mathcal{P}_\lambda} + \frac{1}{T}\langle \mathbb{Q}[\boldsymbol{x}]\rangle_{\mathcal{P}_\lambda} , \tag{10}$$

where

$$\begin{aligned}\mathbb{Q}[\boldsymbol{x}] &= -\mathcal{L}[\boldsymbol{x};\boldsymbol{f}_\lambda] + \mathcal{L}[\boldsymbol{x}^R;\boldsymbol{f}_\kappa] \\ &= \int_0^\tau dt\ \big[\dot{\boldsymbol{x}}\cdot(\boldsymbol{f}_\lambda + \boldsymbol{f}_\mu)/2 - \left(\boldsymbol{f}_\lambda^2 - \boldsymbol{f}_\mu^2\right)/4 \\ &\qquad -T\boldsymbol{\nabla}_{\boldsymbol{x}}\cdot\left(\boldsymbol{f}_\lambda - \boldsymbol{f}_\mu\right)/2\big]\end{aligned} \tag{11}$$

with $\boldsymbol{f}_\lambda = \boldsymbol{f}_{\lambda(t)}(\boldsymbol{x}(t))$ and $\boldsymbol{f}_\mu = \boldsymbol{f}_{\kappa(\tau-t)}(\boldsymbol{x}(t))$. Deriving the second equality in Eq. (11), we made a change of a variable $t \to \tau - t$ for $\mathcal{L}[\boldsymbol{x}^R;\boldsymbol{f}_\kappa]$. The first term in the righthand side of Eq. (10) is equal to $\langle\Delta\mathcal{S}_{sys}\rangle$, which is independent of $\mu(t)$. So it suffices to minimize the average value of $\mathbb{Q}$. The functional $\mathbb{Q}$ will be called a *pseudo-heat* since it reduces to the heat $\mathcal{Q}$ (see Eq. (7)) when $\mu(t) = \lambda(t)$.

The pseudo-heat is minimized at the optimal protocol μ_{op} satisfying $\delta\langle\mathbb{Q}\rangle/\delta\mu(t)|_{\mu_{op}} = 0$. It yields that

$$\left.\left\langle \dot{\boldsymbol{x}}\cdot\frac{\partial \boldsymbol{f}_\mu}{\partial\mu} + \boldsymbol{f}_\mu\cdot\frac{\partial \boldsymbol{f}_\mu}{\partial\mu} + T\boldsymbol{\nabla}_{\boldsymbol{x}}\cdot\frac{\partial \boldsymbol{f}_\mu}{\partial\mu}\right\rangle_{\mathcal{P}_\lambda}\right|_{\mu_{op}} = 0 . \tag{12}$$

It becomes simpler when the force is decomposed as

$$\boldsymbol{f}_\lambda(\boldsymbol{x}) = \boldsymbol{g}(\boldsymbol{x}) + \lambda(t)\boldsymbol{h}(\boldsymbol{x}) \tag{13}$$

with auxiliary force fields $\boldsymbol{g}$ and $\boldsymbol{h}$. Then, the SEP is achieved with the force $\boldsymbol{f}_{\mu^R_{op}(t)}(\boldsymbol{x}) = \boldsymbol{g}(\boldsymbol{x}) + \mu_{op}(\tau - t)\boldsymbol{h}(\boldsymbol{x})$ with the optimal protocol

$$\mu_{op}(t) = -\frac{\langle(\dot{\boldsymbol{x}}(t) + \boldsymbol{g}(\boldsymbol{x}(t)) + T\boldsymbol{\nabla}_{\boldsymbol{x}})\cdot\boldsymbol{h}(\boldsymbol{x}(t))\rangle_{\mathcal{P}_\lambda}}{\langle \boldsymbol{h}(\boldsymbol{x}(t))^2\rangle_{\mathcal{P}_\lambda}} . \tag{14}$$

In the Stratonovich calculus, $\dot{\boldsymbol{x}}(t)\cdot\boldsymbol{F}(\boldsymbol{x}(t))$ should be interpreted as $\lim_{\delta t\to 0}\frac{1}{\delta t}(\boldsymbol{x}(t+\delta t) - \boldsymbol{x}(t))\cdot\frac{1}{2}(\boldsymbol{F}(\boldsymbol{x}(t+\delta t)) + \boldsymbol{F}(\boldsymbol{x}(t)))$ for any vector field $\boldsymbol{F}(\boldsymbol{x})$ [5]. It yields

$$\langle \dot{\boldsymbol{x}}\cdot\boldsymbol{F}(\boldsymbol{x})\rangle_{\mathcal{P}_\lambda} = \langle(\boldsymbol{f}_\lambda + T\boldsymbol{\nabla}_{\boldsymbol{x}})\cdot\boldsymbol{F}(\boldsymbol{x})\rangle_{\mathcal{P}_\lambda} . \tag{15}$$

It further simplifies the optimal condition to the form

$$\mu_{op} = \lambda - 2\frac{\langle\dot{\boldsymbol{x}}\cdot\boldsymbol{h}\rangle_{\mathcal{P}_\lambda}}{\langle\boldsymbol{h}^2\rangle_{\mathcal{P}_\lambda}} = -\lambda - 2\frac{\langle(\boldsymbol{g} + T\boldsymbol{\nabla}_{\boldsymbol{x}})\cdot\boldsymbol{h}\rangle_{\mathcal{P}_\lambda}}{\langle\boldsymbol{h}^2\rangle_{\mathcal{P}_\lambda}} , \tag{16}$$

where function arguments are omitted for simplicity. Hereafter, the average $\langle \cdot \rangle$ is to be taken with respect to $\mathcal{P}_\lambda$ unless stated otherwise. The deviation $(\mu_{op} - \lambda)$ is proportional to the nonequilibrium work done by the nonconservative force. Note that $\delta^2 \langle \mathbb{Q} \rangle / \delta t^2 |_{\mu_{op}} = \langle \boldsymbol{h}^2 \rangle \geq 0$. Hence, $\delta \langle \mathbb{Q} \rangle / \delta t |_{\mu_{op}} = 0$ indeed provides the minimum of the pseudo-heat.

The efficiency of the SEP can be measured by the relative entropy $\mathbb{S}_{op} \equiv \mathbb{S}[\kappa = \mu^R_{op}; \lambda] = \langle \mathcal{S}_{sys} \rangle + \langle \mathbb{Q}_{op} \rangle / T$ with $\mathbb{Q}_{op} = \mathbb{Q}|_{\mu_{op}}$. We define the pseudo-heat dissipation rate $\dot{\mathbb{Q}}_{op}$ as the integrand of Eq. (11) with $\mu = \mu_{op}$. A straightforward calculation shows that

$$\left\langle \dot{\mathbb{Q}}_{op} \right\rangle = \left\langle \dot{\mathcal{Q}} \right\rangle - \frac{1}{4} (\lambda - \mu_{op})^2 \langle \boldsymbol{h}^2 \rangle , \tag{17}$$

where the average heat dissipation rate is given by

$$\langle \dot{\mathcal{Q}} \rangle = \langle \dot{\boldsymbol{x}} \cdot \boldsymbol{f}_\lambda \rangle = \langle (\boldsymbol{f}_\lambda + T \boldsymbol{\nabla}_{\boldsymbol{x}}) \cdot \boldsymbol{f}_\lambda \rangle . \tag{18}$$

It is manifest that $\langle \dot{\mathcal{Q}} \rangle \geq \langle \dot{\mathbb{Q}}_{op} \rangle$.

We will apply the formalism developed to systems with time-dependent and time-independent protocols separately.

Time dependent $\lambda(t)$ *case* – We demonstrate the SEP with two solvable examples. First, consider a harmonic oscillator in one dimension whose stable position is dragged according to a given protocol $\lambda(t)$. The force is given by

$$f_{\lambda(t)}(x) = -k(x - \lambda(t)) \tag{19}$$

with a force constant k. Without loss of generality, we will set $\lambda(0) = 0$. This system describes a bead trapped by an optical tweezer or an electric charge in an electric circuit [20–24].

The probability distribution for $x(t)$ is known exactly (see e.g., Ref. [23]). The particle is assumed to follow the Boltzmann distribution $p_i(x(0)) \propto e^{-kx(0)^2/(2T)}$ at $t = 0$. Then $x(t)$ follows the Gaussian distribution at all t with the mean $\langle x(t) \rangle = \Lambda(t)$ and the variance $\langle (x(t) - \Lambda(t))^2 \rangle = T/k$, where $\Lambda(t) = k \int_0^t ds\ \lambda(s) e^{-k(t-s)}$.

The system corresponds to the case in Eq. (13) with $g(x) = -kx$ and $h(x) = k$. Hence, the optimal protocol from Eq. (16) is given by

$$\mu_{op}(t) = -\lambda(t) + 2\langle x(t) \rangle = -\lambda(t) + 2\Lambda(t) . \tag{20}$$

The probability distribution for $x(t)$ follows the Gaussian distribution with a constant variance at all t. So the system entropy does not change. The pseudo-heat, evaluated from Eqs. (17) and (18) also vanishes while $\langle \dot{\mathcal{Q}} \rangle = k^2 (\lambda - \Lambda)^2$. Therefore the relative entropy $\mathbb{S}_{op}$ vanishes and the SEP is perfect.

Figure 1 illustrates the perfect SEP with a protocol $\lambda(t) = vt$. The initial position $x(0)$ is drawn from the Boltzmann distribution $p_i(x(0)) \sim e^{-kx(0)^2/(2T)}$. In the interval $0 \leq t \leq \tau$, the particle is driven under the protocol $\lambda(t)$. Subsequently, in the interval $\tau \leq t \leq 2\tau$, the particle is driven under the time-reversed protocol $\lambda(2\tau - t)$ or the optimal protocol $\mu_{op}(2\tau - t)$ where $\mu_{op}(t) = -\lambda(t) + 2\Lambda(t) = vt - 2v(1 - e^{-kt})/k$. The SEP is evident under the optimal protocol.



FIG. 1. (Color online) Sample paths of a dragged harmonic oscillator. Thin lines represent paths under λ (black, $0 \leq t \leq \tau$), λ^R (red and dashed, $\tau \leq t \leq 2\tau$), and μ_{op} (blue, $\tau \leq t \leq 2\tau$). Thick lines represent the stable position under corresponding protocols. $k = T = v = 1$ and $\tau = 10$.

As a second example, consider a particle in one dimension subject to a force

$$f_{\lambda(t)}(x) = -\lambda(t) x \tag{21}$$

with a time dependent force constant $\lambda(t)$ [22]. The solution of the Langevin equation is given by $x(t) = x(0) e^{-K(t,0)} + \sqrt{2T} \int_0^t dt' e^{-K(t,t')} \eta(t')$, where $K(t,t') \equiv \int_{t'}^t ds \lambda(s)$. The initial position $x(0)$ is also taken to follow the Boltzmann distribution $p_i(x(0)) \propto e^{-\lambda(0) x(0)^2/(2T)}$. Then, $x(t)$, being the sum of independent Gaussian random variables, follows the Gaussian distribution with the mean $\langle x(t) \rangle = 0$ and the t-dependent variance

$$\langle x(t)^2 \rangle = \frac{T}{\lambda(0)} e^{-2K(t,0)} + 2T \int_0^t dt' e^{-2K(t,t')} . \tag{22}$$

This system corresponds to the case in Eq. (13) with $g(x) = 0$ and $h(x) = -x$. From Eq. (16), we find that the optimal protocol for the SEP is given by

$$\mu_{op}(t) = \frac{2T}{\langle x(t)^2 \rangle} - \lambda(t) . \tag{23}$$

In this case, the probability distribution for $x(t)$ changes its shape. So the system entropy varies with time as (see Eq.(6))

$$\langle \Delta S_{sys} \rangle = \langle S_{sys}(t) \rangle - \langle S_{sys}(0) \rangle = \frac{1}{2} \ln \frac{\langle x(t)^2 \rangle}{\langle x(0)^2 \rangle} . \tag{24}$$

Differentiating it with respect to t and using Eq. (22), one finds that the system entropy increase rate is given by

$$\langle \dot{S}_{sys} \rangle = -\frac{\lambda(t)}{\langle x(t)^2 \rangle} \left(\langle x(t)^2 \rangle - \frac{T}{\lambda(t)} \right) . \tag{25}$$

The pseudo-heat production rate is also calculated using Eqs. (18) and (17). It is given by

$$\langle \dot{\mathbb{Q}}_{op} \rangle = \frac{\lambda(t) T}{\langle x(t)^2 \rangle} \left(\langle x(t)^2 \rangle - \frac{T}{\lambda(t)} \right) . \tag{26}$$

In this case, $\langle \dot{S}_{sys} \rangle + \langle \dot{\mathbb{Q}}_{op} \rangle / T = 0$ while $\langle \dot{S}_{sys} \rangle$ and $\langle \dot{\mathbb{Q}}_{op} \rangle$ are nonzero individually. Therefore the optimal protocol also leads to the perfect SEP.

Time-independent λ case – One can drive a system into a nonequilibrium condition by applying a nonconservative force [6–8]. Then, a total force can be decomposed as

$$\boldsymbol{f}_\lambda(\boldsymbol{x}) = \boldsymbol{f}_c(\boldsymbol{x}) + \lambda \boldsymbol{f}_{nc}(\boldsymbol{x}) , \tag{27}$$

where $\boldsymbol{f}_c(\boldsymbol{x}) = -\boldsymbol{\nabla}_{\boldsymbol{x}} V(\boldsymbol{x})$ denotes a conservative force with a scalar function $V(\boldsymbol{x})$ and $\lambda \boldsymbol{f}_{nc}(\boldsymbol{x})$ denotes a nonconservative force. In this case, a time-independent λ represents the strength of a nonconservative force. In comparison with Eq. (13), $\boldsymbol{g} = \boldsymbol{f}_c$ and $\boldsymbol{h} = \boldsymbol{f}_{nc}$.

Suppose that the system is in the nonequilibrium steady state (NESS) with the probability distribution denoted by $p_{ss}(\boldsymbol{x}) = e^{-\phi_{ss}(\boldsymbol{x})}$. In the NESS, the system entropy is constant while there is an heat dissipation at a constant rate $\langle \dot{\mathcal{Q}} \rangle = \langle \dot{\boldsymbol{x}} \cdot \boldsymbol{f}_\lambda \rangle_{ss} = \lambda \langle \dot{\boldsymbol{x}} \cdot \boldsymbol{f}_{nc} \rangle_{ss}$, where $\langle \cdot \rangle_{ss}$ denotes the average over the NESS [7]. Note that the average rate of the work done by the conservative force is zero, $\langle \dot{\boldsymbol{x}} \cdot \boldsymbol{f}_c \rangle_{ss} = -\frac{d}{dt} \langle V(\boldsymbol{x}(t)) \rangle_{ss} = 0$.

The optimal protocol for the SEP is rephrased as

$$\mu_{op} = -\lambda - 2 \frac{\langle \boldsymbol{f}_c \cdot \boldsymbol{f}_{nc} + T \boldsymbol{\nabla}_{\boldsymbol{x}} \cdot \boldsymbol{f}_{nc} \rangle_{ss}}{\langle \boldsymbol{f}_{nc} \rangle_{ss}^2} . \tag{28}$$

It is constant since the averages are taken over the NESS. From Eqs. (17) and (18), the average pseudo-heat dissipation rate is given by

$$\langle \dot{\mathbb{Q}}_{op} \rangle_{ss} = \frac{\langle \dot{\boldsymbol{x}} \cdot \boldsymbol{f}_{nc} \rangle_{ss}}{\langle \boldsymbol{f}_{nc}^2 \rangle_{ss}} \left(\lambda \langle \boldsymbol{f}_{nc}^2 \rangle_{ss} - \langle \dot{\boldsymbol{x}} \cdot \boldsymbol{f}_{nc} \rangle_{ss} \right) . \tag{29}$$

The pseudo-heat does not vanish in general. Therefore the relative entropy $\mathbb{S}_{op}$ is nonzero and the SEP is imperfect.

Note that the decomposition of a given force $\boldsymbol{f}$ into the form in Eq. (27) is not unique. One can add and subtract a gradient of any scalar function to the conservative and from nonconservative forces, respectively [8]. Making use of the arbitrariness and redefining λ, one can choose

$$\boldsymbol{f}_c(\boldsymbol{x}) = -T \boldsymbol{\nabla}_{\boldsymbol{x}} \phi_{ss}(\boldsymbol{x}) . \tag{30}$$

The NESS $p_{ss}(\boldsymbol{x}) = e^{-\phi_{ss}(\boldsymbol{x})}$ is the steady state solution of the Fokker-Planck equation [5]

$$\frac{\partial p}{\partial t} = \boldsymbol{\nabla} \cdot (-\boldsymbol{f}_\lambda + T \boldsymbol{\nabla}_{\boldsymbol{x}}) \, p . \tag{31}$$

The steady state condition yields that

$$(-(\boldsymbol{\nabla}_{\boldsymbol{x}} \phi_{ss}) + \boldsymbol{\nabla}_{\boldsymbol{x}}) \cdot (\boldsymbol{f}_\lambda + T \boldsymbol{\nabla}_{\boldsymbol{x}} \phi_{ss}) = 0 . \tag{32}$$

If one adopts the choice in Eq. (30), the steady state condition becomes

$$\boldsymbol{f}_c \cdot \boldsymbol{f}_{nc} + T \boldsymbol{\nabla} \cdot \boldsymbol{f}_{nc} = 0 . \tag{33}$$

The converse is also true. If the decomposition in Eq. (27) satisfies Eq. (33), it implies Eq. (30). The steady state condition along this line was mentioned in Ref. [25].

The special choice of $\boldsymbol{f}_c = -T \boldsymbol{\nabla}_{\boldsymbol{x}} \phi_{ss}$ turns out to be extremely useful. The optimal protocol is given by $\mu_{op} = -\lambda$, which means that one needs to reverse the direction of the driving force for the SEP. The pseudo-heat dissipation rate, evaluated directly from Eq. (11) with $\mu = -\lambda$ and Eq. (33), is given by $\mathbb{Q}_{op} = \dot{\boldsymbol{x}} \cdot \boldsymbol{f}_c = -T \frac{d}{dt} \phi_{ss}(\boldsymbol{x}(t))$, whose average value vanishes in the NESS. Therefore, we conclude that a nonequilibrium system driven by a nonconservative force can display the perfect SEP. Precisely speaking, a system being exerted by a force $\boldsymbol{f} = (-T \boldsymbol{\nabla}_{\boldsymbol{x}} \phi_{ss}) + (\boldsymbol{f} + T \boldsymbol{\nabla}_{\boldsymbol{x}} \phi_{ss})$ is equivalent to its dual system being exerted by a dual force $\boldsymbol{f}_{dual} = -T \boldsymbol{\nabla}_{\boldsymbol{x}} \phi_{ss} - (\boldsymbol{f} + T \boldsymbol{\nabla}_{\boldsymbol{x}} \phi_{ss}) = -\boldsymbol{f}(\boldsymbol{x}) - 2T \boldsymbol{\nabla} \phi_{ss}(\boldsymbol{x})$. Both systems have the same NESS and are linked through the time-reversal operation.

Summary and discussion – We have investigated the SEP in nonequilibrium systems driven with a time-dependent force or a nonconservative force. By minimizing the relative entropy in Eq. (10), we have derived the optimal condition for the SEP in Eq. (16). Our study shows that stochastic motions of nonequilibrium system can be reversed under the precise control using the optimal protocol.

The harmonic oscillator system, whose stable position or force constant is varying with time, is shown to display the perfect SEP. It remains uncertain whether the harmonic oscillator is an exceptional case or not. Further studies on even more complicated systems are necessary. Nonequilibrium systems driven by a constant nonconservative force are also shown to display the perfect SEP. As a byproduct, we discover that any driven system has a dual system. They are symmetric under the time reversal and share the same NESS. The duality may shed some light on the general property of the NESS. The pseudo-heat plays an important role in the time-reversibility of nonequilibrium systems. The physical meaning of the pseudo-heat is an open question. The expression in Eq. (29) suggests that the it might be related to the violation of the fluctuation dissipation relation in the NESS [26–28]. It should be scrutinized further. We also propose the the SEP can be used for the test of stochastic thermodynamics in experimental systems.

Acknowledgments – This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MEST) (No. 2012-0005003). We thank Prof. Hyunggyu Park and Prof. Juyeon Yi for helpful discussions.

---

[1] G.I. Taylor, *Low Reynolds Number Flows* (National Committee for Fluid Mechanics Films, Education Development Center, Newton, MA (1966)).
[2] D. J. Pine, J. P. Gollub, J. F. Brady, and A. M. Leshansky, Nature **438**, 997 (2005).
[3] L. Corté, P. M. Chaikin, J. P. Gollub, and D. J. Pine, Nature Physics **4**, 420 (2008).
[4] G. Düring, D. Bartolo, and J. Kurchan, Phys. Rev. E **79**, 030101 (2009).
[5] C. Gardiner, in *Stochastic Methods*, 4th ed. (Springer, Berlin, 2009).
[6] U. Seifert, Phys. Rev. Lett. **95**, 040602 (2005).
[7] T. Hatano and S.-I. Sasa, Phys. Rev. Lett. **86**, 3463 (2001).
[8] C. Kwon, J. D. Noh, and H. Park, Phys. Rev. E **83**, 061145 (2011).
[9] C. Jarzynski, Phys. Rev. Lett. **78**, 2690 (1997).
[10] J. Kurchan, J. Phys. A **31**, 3719 (1998).
[11] G.E. Crooks, Phys. Rev. E **60**, 2721 (1999).
[12] J. Lebowitz and H. Spohn, J. Stat. Phys. **95**, 333 (1999).
[13] D. J. Evans and D. J. Searles, Adv. Phys. **51**, 1529 (2002).
[14] J.D. Noh and J.-M. Park, Phys. Rev. Lett. **108**, (2012).
[15] L. Onsager and S. Machlup, Phys. Rev. **91**, 1505 (1953); S. Machlup and L. Onsager, *ibid.* **91**, 1512 (1953).
[16] A. W. C. Lau and T. C. Lubensky, Phys. Rev. E **76**, 011123 (2007).
[17] T. M. Cover and J. A. Thomas, *Elements of Information Theory*, 2nd ed. (Wiley, 2006).
[18] T. Schmiedl and U. Seifert, Phys. Rev. Lett. **98**, 108301 (2007).
[19] E. Aurell, C. Mejía-Monasterio, and P. Muratore-Ginanneschi, Phys. Rev. Lett. **106**, 250601 (2011).
[20] R. Van Zon, S. Ciliberto, and E.G.D. Cohen, Phys. Rev. Lett. **92**, 130601 (2004).
[21] G. Wang, E. Sevick, E. Mittag, D. Searles, and D. Evans, Phys. Rev. Lett. **89**, (2002).
[22] N. Garnier and S. Ciliberto, Phys. Rev. E **71**, 060101 (2005).
[23] R. Van Zon and E.G.D. Cohen, Phys. Rev. E **69**, 056121 (2004).
[24] A. Imparato, L. Peliti, G. Pesce, G. Rusciano, and A. Sasso, Phys. Rev. E **76**, 050101 (2007).
[25] C. Kwon and P. Ao, Phys. Rev. E 84, 061106 (2011).
[26] T. Harada and S.-I. Sasa, Phys. Rev. Lett. **95**, 130602 (2005).
[27] J. Prost, J.-F. Joanny, and J.M.R. Parrondo, Phys. Rev. Lett. **103**, (2009).
[28] K. Mallick, M. Moshe, and H. Orland, J. Phys. A **44**, 095002 (2011).